

# UCLA

## UCLA Previously Published Works

### Title

Adherence to DASH dietary pattern is inversely associated with osteoarthritis in Americans.

### Permalink

<https://escholarship.org/uc/item/6w3095w0>

### Journal

International journal of food sciences and nutrition, 71(6)

### ISSN

0963-7486

### Authors

Zhang, Yurong  
Francis, Ellen C  
Xia, Tong  
et al.


### Publication Date

2020-09-01

### DOI

10.1080/09637486.2020.1722075

Peer reviewed



## Adherence to DASH dietary pattern is inversely associated with osteoarthritis in Americans

Yurong Zhang, Ellen C. Francis, Tong Xia, Karen Kemper, Joel Williams & Liwei Chen

To cite this article: Yurong Zhang, Ellen C. Francis, Tong Xia, Karen Kemper, Joel Williams & Liwei Chen (2020): Adherence to DASH dietary pattern is inversely associated with osteoarthritis in Americans, International Journal of Food Sciences and Nutrition, DOI: [10.1080/09637486.2020.1722075](https://doi.org/10.1080/09637486.2020.1722075)

To link to this article: <https://doi.org/10.1080/09637486.2020.1722075>



Published online: 04 Feb 2020.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)



RESEARCH ARTICLE



## Adherence to DASH dietary pattern is inversely associated with osteoarthritis in Americans

Yurong Zhang<sup>a</sup>, Ellen C. Francis<sup>b</sup>, Tong Xia<sup>c</sup>, Karen Kemper<sup>b</sup>, Joel Williams<sup>b</sup> and Liwei Chen<sup>c</sup>

<sup>a</sup>Department of Neurology, The First Affiliated Hospital of Xi'an Jiaotong University, Shaanxi, China; <sup>b</sup>Department of Public Health Sciences, Clemson University, Clemson, SC, USA; <sup>c</sup>Fielding School of Public Health, University of California, Los Angeles, CA, USA

### ABSTRACT

Osteoarthritis (OA) is one of the most common diseases that cause disability among older adults. The objective of this study was to assess the association between adherence to the Dietary Approaches to Stop Hypertension (DASH) and OA in American adults. This study included adults ( $\geq$  aged 20 years) who participated in the National Health and Examination Survey (NHANES) 2007–2016 in the United States. Adherence to the DASH score was calculated from 8 food groups. Higher scores indicate better adherence to the DASH dietary pattern. Among the 21,901 participants included in this study, 10.26% reported having OA. Results of our multivariable logistic regression indicated a statistically significant inverse association between DASH score tertiles and OA. The adjusted ORs (95% CI) were 1.00 (ref), 0.89 (0.72; 1.10), and 0.78 (0.60; 1.00) across increasing DASH score tertiles ( $P$  for trend = 0.045). In this representative sample of American adults, greater adherence to the DASH dietary pattern was associated with lower likelihood of having OA.

### ARTICLE HISTORY

Received 1 October 2019  
Revised 20 January 2020  
Accepted 23 January 2020

### KEYWORDS

DASH diet; osteoarthritis; adults; nutrition; degenerative joint disease

## Introduction

Osteoarthritis (OA) is the most prevalent degenerative joint disease and the fastest growing cause of disability and pain in the United State (US). Among US adults 20 years and older, the prevalence of OA increased from 6.6% in 1999 to 14.3% in 2014 (Park et al. 2018). The number of people with OA is expected to increase in the US due in part to the obesity epidemic and aging of the population. Globally, approximately 18% of women and 9.6% of men aged over 60 years have symptomatic OA, and by 2050, OA is projected to affect 130 million people (Conaghan et al. 2015).

OA is one of the most challenging joint diseases for clinicians. Besides symptom management and surgical approaches, there is no cure for OA (Zhang et al. 2016). Recently, dietary factors have received considerable interest due to their potential roles in OA prevention. Most previous studies on dietary intake and OA only focussed on an individual nutrient. For example, vitamin D (McAlindon et al. 1996; Lane et al. 1999) and vitamin K (Misra et al. 2013; Shea et al. 2015) play a physiological role in bone and cartilage mineralization, and have been associated with risk of OA in several prospective cohort studies.

However, randomised controlled trials on vitamin D, K, or E supplementation have failed to show success in improving the symptoms and progression of OA (Neogi et al. 2008; Arden et al. 2016; Jin et al. 2016; Diao et al. 2017). One explanation could be that individual nutrients only have a limited effect, and therefore the impact may be difficult to detect in single-nutrient supplementation trials. To gain a better understanding of dietary factors in relation to OA development or progression, studies investigating overall dietary quality/pattern are needed. However, such studies are sparse, and to our best knowledge, only two studies have examined the association between an individual's dietary pattern and OA risk (Veronese et al. 2017; Veronese et al. 2019).

The Dietary Approaches to Stop Hypertension (DASH) dietary pattern, which was originally developed to reduce blood pressure, is low in red meat, sweets and oils; and rich in fruits, vegetables, low-fat dairy products, whole grains, fish, and nuts (Appel et al. 1997). In the recent Dietary Guidelines for Americans (2015–2020), the United States Department of Agriculture recommended the DASH dietary pattern as a healthy eating plan for Americans based on

growing evidence that adherence to the DASH dietary pattern had beneficial effects on preventing type II diabetes, cardiovascular diseases (CVD), metabolic syndrome, and Alzheimer's disease, in addition to hypertension (Asemi et al. 2013; Nilsson et al. 2019; Pickworth et al. 2019; Samadi et al. 2019). A recent randomised cross-over study among children showed that the DASH diet significantly reduced the circulating levels of serum high-sensitivity C-reactive protein (hs-CRP: a biomarker of chronic inflammation), independent of changes in body weight and lipid profile (Saneei et al. 2014). Therefore, we hypothesised that adherence to the DASH dietary pattern would be associated with lower risk of OA. The objective of this study was to examine the association of adherence to the DASH dietary pattern with OA in American adults.

## Methods

### Study population

The study population included participants from the National Health and Nutrition Examination Survey (NHANES) 2007–2016. The NHANES is a nationally representative cross-sectional survey of the non-institutionalized US population aged 2 years or older with data collected in 2-year cycles. During each cycle, the NHANES was conducted based on a stratified multi-stage probability sampling design and included two components: a household interview and a health examination. The health examination component consisted of medical, dental, and physiological measurements, as well as laboratory tests administered by trained medical personnel in a fully equipped mobile examination centre (MEC). Additional information on the design and procedures of the NHANES are available at the Centre for Disease Control and Prevention (CDC) website. For this study, we included individuals who were aged 20 years or older and participated in the NHANES 2007–2016 survey cycles, which includes the most recent dietary data released for public use. We excluded participants who declined or had missing data on arthritis information, those who had no or invalidated data on dietary intake, had total energy intake  $<500$  or  $>8000$  kcal/day, or who reported being on a special diet for medical reasons or to lose weight. The final sample size included in this analysis was 21,901. The NHANES survey was approved by the Research Ethics Review Board of the National Centre for Health Statistics (NCHS) and conducted following the rules of the Declaration of Helsinki.

### Assessment of osteoarthritis (OA)


OA was assessed using the data from the medical conditions questionnaire which was collected during the NHANES household interview. Participants were classified as having OA if they reported ever being told they had arthritis by a health care provider and indicated that it was osteoarthritis.

### Adherence to dietary approaches to stop hypertension (DASH)

Dietary intake was assessed by two 24-h dietary recalls administered by trained interviewers. We constructed the DASH score based on intakes of 8 DASH targeted food groups/items (i.e. whole grains, vegetables, fruits, dairy products, red meats, nuts/seeds/legumes, sweets, and sodium) using the method developed by Fung et al. (Fung et al. 2008). Briefly, we ranked study participants according to their intake of each individual food group/item and classified them into quintiles. We assigned a score of 1 for individuals in the 1st quintile and 5 for individuals in the 5<sup>th</sup> quintile for whole grains, vegetables, fruits, dairy products, nuts/seeds/legumes because high intake is desired for these foods. For red meats, sweets, and sodium, we inverted scores assigning 5 for the 1st quintile and 1 for the 5th quintile because low intake is desired for these food groups/items. We then summed the score of each food group/item to generate an overall DASH adherence score ranging from 8 (lowest adherence) to 40 (best adherence).

### Covariates

We considered the following variables as potential confounders in examining the association between adherence to the DASH dietary pattern and OA. Age, sex, race/ethnicity, annual household income, education, and smoking, which were self-reported during the household interviews. Alcohol intake was assessed by 24-h dietary recalls. Physical activity (PA) was measured using a structured questionnaire and calculated as the sum of the minutes spent being physically active for commuting, recreation, and work on average for each day. Adequate PA was defined as time of all PA  $\geq 150$  minutes per week. Height, weight, and waist circumference were directly measured by trained personnel in the MEC. Body mass index (BMI) was calculated by weight (kg) divided height squared ( $\text{m}^2$ ).



**Figure 1.** Prevalence (%) of osteoarthritis in NHANES 2007–2016 by age group.

### Statistical analysis

Descriptive data on study participants' characteristics were expressed as weighted means (standard error) for continuous variables or weighted percent for categorical variables if not mentioned otherwise. Student's *t*-test and  $\chi^2$  test were applied to compare continuous variables and categorical variables, respectively. The prevalence of OA was estimated in all participants and by age groups (20–29.9, 30–39.9, 40–49.9, 50–59.9, 60–69.9,  $\geq 70$  years). Logistic regressions were performed to estimate the association of DASH score with OA. The DASH score was analysed both as a categorical variable (tertiles) and a continuous variable. Because age is a strong risk factor for OA, all models were adjusted for age. Other covariates controlled for in the multivariable regression models included sex (female versus male), race/ethnicity (non-Hispanic White, non-Hispanic Black, or Mexican American/other Hispanics), annual household income ( $< \$20,000$ ,  $\$20,000$ – $49,999$ ,  $\$45,000$ – $\$74,999$ ,  $\geq \$75,000$ ), education (less than high school, high school, some college, college graduate or above), smoking (current, former, or never), alcohol intake (g/day), PA (adequate vs. non-adequate), and BMI (normal weight: BMI  $< 25$ ; overweight: BMI 25–29.9;

obese: BMI  $\geq 30$  kg/m<sup>2</sup>). Additionally, we performed stratified analyses by gender (males and females), BMI (normal weight, overweight and obese), and smoking status (smokers and never smokers). All analyses considered the complex survey design including stratification, clustering, and weighting to account for the oversampling of subgroups, unit non-response and non-coverage in the NHANES. All the statistical analyses were performed using SAS software, version 9.4 (SAS Institute, Cary, NC).

### Results

Among the 21,901 participants included in this study, the weighted prevalence of OA was 10.26% (0.35). The prevalence of OA increased substantially with age: 0.78% in age group 20–29.9 years, 2.61% in 30–39.9 years, 5.67% in 40–49.9 years, 11.69% in 50–59.9 years, 19.73% in 60–69.9 years, and 27.13% in  $\geq 70$  years, respectively (Figure 1).

The median DASH score was 24.0 with a range from 9.0 to 39. The median (min, max) DASH score in each tertile was 19 (9, 21), 24 (22, 25) and 28 (26, 39), respectively. Characteristics of the study participants in total and by DASH score tertiles are presented in Table 1. Compared to individuals in the

**Table 1.** Characteristics of study participants, NHANES 2007–2016.

Characteristics	DASH score				<i>p</i> Values
	All ( <i>N</i> = 21,901)	Tertile 1 ( <i>N</i> = 6957)	Tertile 2 ( <i>N</i> = 6917)	Tertile 3 ( <i>N</i> = 8027)	
DASH score, median (range)	24 (9, 39)	19 (9, 21)	24 (22, 25)	28 (26, 39)	
Age (years) <sup>a</sup>	46.80 (0.25)	42.11 (0.32)	47.16 (0.33)	51.42 (0.37)	<0.0001
Female, %	49.94 (0.38)	61.27 (0.69)	47.81 (1.02)	39.99 (0.65)	<0.0001
Race, %					<0.0001
Non-Hispanic White	67.16 (1.62)	70.96 (1.76)	66.23 (1.74)	64.05 (1.69)	
Non-Hispanic Black	11.29 (0.83)	12.75 (1.12)	12.03 (0.94)	9.04 (0.63)	
Hispanic	14.33 (1.12)	11.28 (1.06)	14.78 (1.21)	17.13 (1.27)	
Multiracial or other	7.21 (0.45)	5.01 (0.37)	6.96 (0.55)	9.78 (0.26)	
Education Status, %					<0.0001
Less than high school	17.03 (0.71)	18.00 (0.92)	16.82 (0.84)	16.20 (0.78)	
High school	23.04 (0.58)	27.47 (0.82)	23.71 (0.81)	17.69 (0.55)	
Some college	31.05 (0.55)	33.89 (0.82)	31.69 (0.80)	27.42 (0.89)	
College graduate or above	28.89 (1.01)	20.64 (1.23)	27.77 (1.16)	38.69 (1.17)	
Income, %					<0.0001
<20,000	15.11 (0.63)	16.15 (0.87)	14.93 (0.68)	14.16 (0.68)	
20,000–49,999	26.53 (0.68)	27.31 (0.92)	25.70 (0.85)	26.50 (0.80)	
45,000–74,999	21.83 (0.66)	23.30 (0.86)	21.61 (0.91)	20.44 (0.94)	
75,000+	36.53 (1.17)	33.23 (1.33)	37.75 (1.22)	38.90 (1.42)	
Smoking status, %					<0.0001
Current	21.38 (0.56)	33.88 (0.94)	9.44 (0.46)	9.44 (0.46)	
Former	23.62 (0.50)	21.46 (0.78)	24.89 (0.79)	24.89 (0.79)	
Never	55.0 (0.69)	44.67 (1.01)	65.67 (0.90)	65.67 (0.90)	<0.0001
Body Mass Index (kg/m <sup>2</sup> ) <sup>a</sup>	28.51 (0.09)	29.46 (0.12)	28.70 (0.13)	27.32 (0.12)	<0.0001
Waist circumference (cm) <sup>a</sup>	98.10 (0.23)	100.55 (0.31)	98.44 (0.36)	95.11 (0.29)	<0.0001
Dietary intake <sup>a</sup>					
Alcohol drinking, g/day	172.58 (6.13)	289.78 (13.92)	136.32 (6.61)	83.04 (4.02)	<0.0001
Sodium, mg/day	3540 (15.60)	3973 (29.45)	3503 (29.22)	3156 (19.64)	<0.0001
Potassium, mg/day	2694.33 (14.50)	2511.25 (20.65)	2644.15 (21.39)	2936.37 (17.07)	<0.0001
Magnesium, mg/day	302.05 (1.92)	278.01 (2.38)	294.48 (2.46)	334.58 (2.47)	<0.0001
Saturated fat, g/day	26.52 (0.19)	29.64 (0.28)	26.79 (0.28)	23.41 (0.24)	<0.0001
Fibre, g/day	17.09 (0.14)	13.91 (0.13)	16.63 (0.18)	20.91 (0.19)	<0.0001
Vitamin D, mcg/day	4.68 (0.04)	3.74 (0.07)	4.73 (0.09)	5.63 (0.06)	<0.0001
Vitamin K, mcg/day	112.4 (3.04)	96.7 (7.44)	108.2 (2.38)	134.1 (2.66)	<0.0001
Vitamin E, mg/day	8.48 (0.08)	7.94 (0.09)	8.27 (0.10)	9.26 (0.13)	<0.0001
Vitamin C, mg/g	82.18 (1.16)	57.79 (1.47)	79.5 (1.35)	110.5 (1.40)	<0.0001

<sup>a</sup>Mean (SE).

highest DASH score tertile (tertile 3), those in the lowest DASH score tertile (tertile 1) were younger, had a greater BMI and waist circumference, were more likely to be female, non-Hispanic white (70.96% versus 64.05%), to be a current smoker, have an annual household income <\$20,000, and were less likely to have a college degree or above. Individuals in the lowest DASH score tertile had lower intakes of dietary fibre, potassium, magnesium, vitamin D, K, E, and C, but higher intakes of alcohol, sodium, and saturated fat than individuals in the highest tertile.

The DASH score was inversely associated with OA. The age-adjusted ORs (95% CIs) of OA across the DASH score tertiles were 1.00 (reference), 0.86 (0.72; 1.05), and 0.76 (0.63; 0.92), respectively (*P* for trend: 0.0005) (Table 2). The ORs of OA were similar following additional adjustment for gender, race/ethnicity, education, household income, smoking, alcohol drinking, and PA ((1.00 (reference), 0.85 (0.69; 1.04), and 0.74 (0.59; 0.93) across the DASH tertiles) (*P* for trend: 0.0003)). Further adjustment for BMI did not substantially change the results ((1.00 (reference); 0.89 (0.72; 1.10), and 0.78 (0.60; 1.00) across the DASH

tertiles)), however, the statistical significance of the linear trend across tertiles became borderline (*p* = 0.045). When modelling the DASH score continuously, a similar inverse association was observed in fully adjusted models (Model 3 in Table 2). A one unit increase in the DASH score was associated with a 3% lower odds (95% CI: 1–6%; *p* = 0.007) of having OA.

In the stratified analyses, the DASH-OA association was slightly stronger among obese individuals or smokers (combined current smokers and former smokers since there was no difference between these two groups regarding the ORs), but the tests for interactions were all statistically insignificant (Table 3).

## Discussion

In this large study among a US representative sample, we found that adherence to the DASH dietary pattern was inversely associated with OA. Individuals with better adherence to the DASH dietary pattern had lower odds of having OA: individuals in the highest DASH adherence group (tertile 3) had a median



**Table 2.** Unadjusted and adjusted associations between DASH score and osteoarthritis, NHANES 2007–2016.

DASH Score	Sample Size (n)	Age-adjusted Model OR (95% CI)	p Value (trend)	Model 2 OR (95% CI)	p Value (trend)	Model 3 (fully adjusted) OR (95% CI)	p Value (trend)
Categorical							
Tertile 1	6957	Reference		Reference		Reference	
Tertile 2	6917	0.86 (0.72, 1.05)		0.85 (0.69, 1.04)		0.89 (0.72, 1.10)	
Tertile 3	8027	0.76 (0.63, 0.92)	0.0005	0.74 (0.59, 0.93)	0.0003	0.78 (0.60, 1.00)	0.045
Continuous							
DASH score	21,901	0.97 (0.96, 0.99)	0.0004	0.96 (0.94, 0.98)	0.0003	0.97 (0.94, 0.99)	0.007

Model 1: Age.

Model 2: Model 1+ gender, race/ethnicity, education, household income, smoking, alcohol drinking, and physical activity.

Model 3: Model 2+ body mass index.

**Table 3.** Adjusted associations between DASH score and osteoarthritis by gender, smoking status and BMI, NHANES 2007–2016.

	DASH score (continuous)	Sample size (n)	Multivariable-adjusted Model OR (95% CI)	p Value	p Value (interaction)
All		21,901	0.97 (0.94, 0.99)	0.007	
By Gender	Males	13,670	0.97 (0.93, 1.00)	0.51	0.78
	Females	10,945	0.97 (0.94, 1.00)	0.50	
By BMI	Normal weight	6777	0.98 (0.95, 1.02)	0.30	0.39
	Overweight	7306	0.98 (0.94, 1.02)	0.39	
	Obese	7663	0.94 (0.91, 0.98)	0.006	
Smoking	Current and former	9903	0.98 (0.96, 1.02)	0.16	0.11
	Never	12,038	0.95 (0.92, 0.99)	0.009	

Model adjusted for age, gender, race/ethnicity, education, household income, smoking, alcohol drinking, physical activity, and body mass index

DASH score of 28 and a 22% lower likelihood of having OA compared to individuals in the lowest DASH adherence group (tertile 1) who had a median DASH score of 19. A one unit increase in the DASH score was associated with a 3% lower of odds of having OA, independent of sociodemographic and lifestyle factors such as age, education, race/ethnicity, household income, smoking, alcohol consumption, PA and BMI. To our knowledge, this is the first study to investigate the association between adherence to the DASH dietary pattern and OA in American adults. Our results are in line with the previous two studies that investigated the association of OA with adherence to a Mediterranean diet (Veronese et al. 2017) or the Dietary Inflammatory Index (Veronese et al. 2019). Our findings in conjunction with previous studies provide support that overall dietary quality/pattern may play an important role in OA development and should be considered in future interventions.

OA may be amenable to early prevention and treatment, and therefore modifiable factors should be identified. Diet is one of the modifiable risk factors of OA (Georgiev and Angelov 2019). Although the DASH diet was initially designed for lowering the blood pressure among individuals with hypertension, numerous studies have shown that this diet also has other metabolic benefits in different populations (Asghari et al. 2016; Razavi Zade et al. 2016; Kang et al. 2018; Tiong et al. 2018; Akhlaghi 2019). Our findings of an inverse association between adherence to the DASH dietary pattern and OA add new evidence to the existing

literature on potential benefits of the DASH diet for bone and joint health.

Although the association between adherence to the DASH dietary pattern and OA are novel, it is biologically plausible. First the DASH diet is rich in key nutrients required by the bones and tissues of synovial joints, including vitamin D, K, calcium, and magnesium. Compared to adults consuming a typical American diet, continual consumption of the DASH diet over a 30-day period has been associated with a reduction in bone turnover markers and improvement in calcium metabolism (Doyle and Cashman 2004). Deficient dietary consumption of calcium leads to lower bone and cartilage mineral content and bone mineral density, and long-term deficiency can lead to rickets, osteomalacia and osteoporosis (Dermience et al. 2015). Magnesium deficiency may play a role in OA onset and progression. Lower magnesium intake was associated with greater knee pain in individuals with radiographic knee OA (Shmagel et al. 2018). Second, the DASH diet may be beneficial to OA development through an anti-inflammatory pathway (Samadi et al. 2019). There is emerging evidence that the development and progression of OA is associated with low-grade systemic inflammation. A systematic review of 32 studies found that serum levels of hs-CRP were statistically significantly higher in OA patients than in controls (Jin et al. 2015). Inflammation may increase the risk of OA through increased cartilage degeneration and osteophytosis (Kozijn et al. 2019). In a cross-over trial the DASH

diet has been associated with lower levels of hs-CRP/CRP (King et al. 2007). The DASH diet is comprised of many nutrients that may modulate both acute and chronic inflammation (Calder et al. 2009). For example, high dietary saturated fatty acids (Cruz-Teno et al. 2012) have been associated with pro-inflammatory effects, whereas dietary fibre (King et al. 2007) and flavonoids (Saneei et al. 2014) have been associated with anti-inflammatory effects. The DASH diet is rich in flavonoid and fibre but lower in saturated fatty acids (Appel et al. 1997). However, this study was not designed to identify the relationship of each nutrient or food group with OA. Future work is needed to investigate whether this association is causal and to elucidate the underlying mechanisms.

This study has several notable strengths. First, this study included a large nationally representative sample with a wide age range (e.g. 20–80 years). Second, the dietary intakes in NHANES were assessed by two 24-h dietary recalls that were administered by trained research staff and included multiple prompts to reduce the likelihood of missing food or drinks consumed by the participants. Lastly, NHANES had extensive measures of sociodemographic and lifestyle factors, as well as directly measured anthropometric data (e.g. weight, height, and waist circumferences) which allowed us to control for most potential confounders. However, the findings of the current study should be considered with several limitations. NHANES is designed as a cross-sectional study and observational in nature; therefore, causality cannot be determined and residual confounding cannot be completely rule out. Additionally, OA was self-reported and likely to introduce a bias. However, the questionnaire has been widely used in studies to assess self-reported OA (Kojima et al. 2017; Hackney et al. 2019; Parekh et al. 2019).

In conclusion, adherence to the DASH dietary pattern was associated with lower likelihood of having OA among US adults. This is particularly relevant given the increasing prevalence of OA and the cost of treating such chronic conditions. Future studies with prospective designs and more accurate assessment of OA, or clinical trials are needed to confirm the results from current study.

### Authors' contributions

YZ: study conception and design, literature search, interpretation of analyses, drafting the manuscript and major editing; ECF: study conception and design, obtain data, and statistical analysis; XT: statistical

analysis and editing; JW: major editing; KK: major editing; LC: study conception and design, literature search, data obtain and statistical analysis, interpretation of analyses, major editing; All authors read and approved the final manuscript. LC ([cliwei86@ucla.edu](mailto:cliwei86@ucla.edu)) had full access to all the data in the study and took responsibility for the integrity of the data and the accuracy of the data analysis.

### Disclosure statement

The authors declare that they have no conflict of interest.

### References

- Akhlaghi M. 2019. Dietary Approaches to Stop Hypertension (DASH): potential mechanisms of action against risk factors of the metabolic syndrome. *Nutr Res Rev.* 30:1–18.
- Appel LJ, Moore TJ, Obarzanek E, Vollmer WM, Svetkey LP, Sacks FM, Bray GA, Vogt TM, Cutler JA, Windhauser MM, et al. 1997. A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group. *N Engl J Med.* 336:1117–1124.
- Arden NK, Cro S, Sheard S, Doré CJ, Bara A, Tebbs SA, Hunter DJ, James S, Cooper C, O'Neill TW, et al. 2016. The effect of vitamin D supplementation on knee osteoarthritis, the VIDEO study: a randomised controlled trial. *Osteoarthritis Cartilage.* 24:1858–1866.
- Asemi Z, Samimi M, Tabassi Z, Sabihi SS, Esmailzadeh A. 2013. A randomized controlled clinical trial investigating the effect of DASH diet on insulin resistance, inflammation, and oxidative stress in gestational diabetes. *Nutrition.* 29:619–624.
- Asghari G, Yuzbashian E, Mirmiran P, Hooshmand F, Najafi R, Azizi F. 2016. Dietary Approaches to Stop Hypertension (DASH) dietary pattern is associated with reduced incidence of metabolic syndrome in children and adolescents. *J Pediatr.* 174:178–184.e1.
- Calder PC, Albers R, Antoine JM, Blum S, Bourdet-Sicard R, Ferns GA, Folkerts G, Friedmann PS, Frost GS, Guarner F, et al. 2009. Inflammatory disease processes and interactions with nutrition. *Br J Nutr.* 101:S1–S45.
- Conaghan PG, Porcheret M, Kingsbury SR, Gammon A, Soni A, Hurley M, Rayman MP, Barlow J, Hull RG, Cumming J, et al. 2015. Impact and therapy of osteoarthritis: the Arthritis Care OA Nation 2012 survey. *Clin Rheumatol.* 34:1581–1588.
- Cruz-Teno C, Perez-Martinez P, Delgado-Lista J, Yubero-Serrano EM, Garcia-Rios A, Marin C, Gomez P, Jimenez-Gomez Y, Camargo A, Rodriguez-Cantalejo F, et al. 2012. Dietary fat modifies the postprandial inflammatory state in subjects with metabolic syndrome: the LIPGENE study. *Mol Nutr Food Res.* 56:854–865.
- Dermience M, Lognay G, Mathieu F, Goyens P. 2015. Effects of thirty elements on bone metabolism. *J Trace Elem Med Biol.* 32:86–106.
- Diao N, Yang B, Yu F. 2017. Effect of vitamin D supplementation on knee osteoarthritis: a systematic review and



- meta-analysis of randomized clinical trials. *Clin Biochem.* 50:1312–1316.
- Doyle L, Cashman KD. 2004. The DASH diet may have beneficial effects on bone health. *Nutr Rev.* 62:215–220.
- Fung TT, Chiuve SE, McCullough ML, Rexrode KM, Logroscino G, Hu FB. 2008. Adherence to a DASH-style diet and risk of coronary heart disease and stroke in women. *Arch Intern Med.* 168:713–720.
- Georgiev T, Angelov AK. 2019. Modifiable risk factors in knee osteoarthritis: treatment implications. *Rheumatol Int.* 39:1145–1157.
- Hackney AJ, Klindinst NJ, Resnick B, Johantgen M. 2019. Association of systemic inflammation and fatigue in osteoarthritis: 2007–2010 National Health and Nutrition Examination Survey. *Biol Res Nurs.* 21:532–543.
- Jin X, Beguerie JR, Zhang W, Blizzard L, Otahal P, Jones G, Ding C. 2015. Circulating C reactive protein in osteoarthritis: a systematic review and meta-analysis. *Ann Rheum Dis.* 74:703–710.
- Jin X, Jones G, Cicuttini F, Wluka A, Zhu Z, Han W, Antony B, Wang X, Winzenberg T, Blizzard L, et al. 2016. Effect of vitamin D supplementation on tibial cartilage volume and knee pain among patients with symptomatic knee osteoarthritis: a randomized clinical trial. *JAMA.* 315:1005–1013.
- Kang SH, Cho KH, Do JY. 2018. Association between the modified dietary approaches to stop hypertension and metabolic syndrome in postmenopausal women without diabetes. *Metab Syndr Relat Disord.* 16:282–289.
- King DE, Egan BM, Woolson RF, Mainous AG 3rd, Al-Solaiman Y, Jesri A. 2007. Effect of a high-fiber diet vs a fiber-supplemented diet on C-reactive protein level. *Arch Intern Med.* 167:502–506.
- Kojima N, Kim M, Saito K, Yoshida Y, Hirano H, Obuchi S, Shimada H, Suzuki T, Kim H. 2017. Predictors of self-reported knee osteoarthritis in community-dwelling older women in Japan: a cross-sectional and longitudinal cohort study. *Arch Gerontol Geriatr.* 73:125–132.
- Kozijn AE, Tartjiono MT, Ravipati S, van der Ham F, Barrett DA, Mastbergen SC, Korthagen NM, Lafeber F, Zuurmond AM, Bobeldijk I, et al. 2019. Human C-reactive protein aggravates osteoarthritis development in mice on a high-fat diet. *Osteoarthritis Cartilage.* 27:118–128.
- Lane NE, Gore LR, Cummings SR, Hochberg MC, Scott JC, Williams EN, Nevitt MC; for the Study of Osteoporotic Fractures Research Group. 1999. Serum vitamin D levels and incident changes of radiographic hip osteoarthritis: a longitudinal study. *Study of Osteoporotic Fractures Research Group. Arthritis Rheum.* 42:854–860.
- McAlindon TE, Jacques P, Zhang Y, Hannan MT, Aliabadi P, Weissman B, Rush D, Levy D, Felson DT. 1996. Do antioxidant micronutrients protect against the development and progression of knee osteoarthritis? *Arthritis Rheum.* 39:648–656.
- Misra D, Booth SL, Tolstykh I, Felson DT, Nevitt MC, Lewis CE, Torner J, Neogi T. 2013. Vitamin K deficiency is associated with incident knee osteoarthritis. *Am J Med.* 126:243–248.
- Neogi T, Felson DT, Sarno R, Booth SL. 2008. Vitamin K in hand osteoarthritis: results from a randomised clinical trial. *Ann Rheum Dis.* 67:1570–1573.
- Nilsson A, Halvardsson P, Kadi F. 2019. Adherence to DASH-style dietary pattern impacts on adiponectin and clustered metabolic risk in older women. *Nutrients.* 11: 805.
- Parekh SM, Fernandes GS, Moses JP, Fuller CW, Scammell BE, Batt ME, Zhang W, Doherty M. 2019. Risk factors for knee osteoarthritis in retired professional footballers: a cross-sectional study. *Clin J Sport Med.*
- Park J, Mendy A, Vieira ER. 2018. Various types of arthritis in the United States: prevalence and age-related trends from 1999 to 2014. *Am J Public Health.* 108:256–258.
- Pickworth CK, Deichert DA, Corroon J, Bradley RD. 2019. Randomized controlled trials investigating the relationship between dietary pattern and high-sensitivity C-reactive protein: a systematic review. *Nutr Rev.* 77:363–375.
- Razavi Zade M, Telkabadi MH, Bahmani F, Salehi B, Farshbaf S, Asemi Z. 2016. The effects of DASH diet on weight loss and metabolic status in adults with non-alcoholic fatty liver disease: a randomized clinical trial. *Liver Int.* 36:563–571.
- Samadi M, Moradi S, Moradinazar M, Mostafai R, Pasdar Y. 2019. Dietary pattern in relation to the risk of Alzheimer's disease: a systematic review. *Neurol Sci.* 40: 2031–2043.
- Saneei P, Hashemipour M, Kelishadi R, Esmailzadeh A. 2014. The Dietary Approaches to Stop Hypertension (DASH) diet affects inflammation in childhood metabolic syndrome: a randomized cross-over clinical trial. *Ann Nutr Metab.* 64:20–27.
- Shea MK, Kritchevsky SB, Hsu FC, Nevitt M, Booth SL, Kwok CK, McAlindon TE, Vermeer C, Drummen N, Harris TB, et al. 2015. The association between vitamin K status and knee osteoarthritis features in older adults: the Health, Aging and Body Composition Study. *Osteoarthritis Cartilage.* 23:370–378.
- Shmagel A, Onizuka N, Langsetmo L, Vo T, Foley R, Ensrud K, Valen P. 2018. Low magnesium intake is associated with increased knee pain in subjects with radiographic knee osteoarthritis: data from the Osteoarthritis Initiative. *Osteoarthritis Cartilage.* 26:651–658.
- Tiong XT, Nursara Shahirah A, Pun VC, Wong KY, Fong AYY, Sy RG, Castillo-Carandang NT, Nang EEK, Woodward M, van Dam RM, et al. 2018. The association of the dietary approach to stop hypertension (DASH) diet with blood pressure, glucose and lipid profiles in Malaysian and Philippines populations. *Nutr Metab Cardiovasc Dis.* 28:856–863.
- Veronese N, Shivappa N, Stubbs B, Smith T, Hebert JR, Cooper C, Guglielmi G, Reginster JY, Rizzoli R, Maggi S. 2019. The relationship between the dietary inflammatory index and prevalence of radiographic symptomatic osteoarthritis: data from the Osteoarthritis Initiative. *Eur J Nutr.* 58:253–260.
- Veronese N, Stubbs B, Noale M, Solmi M, Luchini C, Smith TO, Cooper C, Guglielmi G, Reginster JY, Rizzoli R, et al. 2017. Adherence to a Mediterranean diet is associated with lower prevalence of osteoarthritis: data from the osteoarthritis initiative. *Clin Nutr.* 36:1609–1614.
- Zhang W, Ouyang H, Dass CR, Xu J. 2016. Current research on pharmacologic and regenerative therapies for osteoarthritis. *Bone Res.* 4:15040.